arXiv:1606.00267v1 [math.NT] 1 Jun 2016

A NOTE ON CATALAN NUMBERS ASSOCIATED WITH p-ADIC INTEGRAL ON $\mathbb{Z}_p$

TAEKYUN KIM

Abstract. In this paper, we study Catalan numbers which can be represented by the p-adic integral on $\mathbb{Z}_p$ and we investigate some properties and formulae related to Catalan numbers and special numbers.

1. Introduction

Let p be a f ixed odd prime number. Throughout this paper, $\mathbb{Z}_p$,$\mathbb{Q}_p$ and $\mathbb{C}_p$ will denote the ring of p-adic integers, the field of p-adic numbers and the completion of the algebraic closure of $\mathbb{Q}_p$. The p-adic norm $|\cdot|_p$ is normalized as $|p|_p = \frac{1}{p}$. As is well known, the Euler numbers are defined by the generating function to be

$$\frac{2}{e^t+1}e^{xt} = \sum_{n=0}^{\infty} E_n(x)\frac{t^n}{n!}, \quad (\text{see } [1-13]). \tag{1.1}$$

When $x = 0$, $E_n = E_n(0)$ are called the Euler numbers. From (1.1), we note that

$$E_n(x) = \sum_{l=0}^{n} \binom{n}{l} x^{n-l} E_l. \tag{1.2}$$

Recently, λ-Changhee polynomials are defined by the generating function to be

$$\frac{2}{(1+t)^{\lambda}+1}(1+t)^{\lambda x} = \sum_{n=0}^{\infty} Ch_{n,\lambda}(x)\frac{t^n}{n!}, \quad \text{where } \lambda \in \mathbb{Z}_p \quad (\text{see } [5]). \tag{1.3}$$

2010 *Mathematics Subject Classification.* 11B68, 11S80.

Key words and phrases. Catalan numbers, fermionic p-adic integral on $\mathbb{Z}_p$.

.

By replacing t by $e^{\frac{1}{\lambda}t}-1$, we get

$$\begin{aligned}\frac{2}{e^t+1}e^{xt} &= \sum_{n=0}^{\infty} Ch_{n,\lambda}(x)\frac{1}{n!}\left(e^{\frac{1}{\lambda}t}-1\right)^n \\ &= \sum_{n=0}^{\infty} Ch_{n,\lambda}(x)\sum_{m=n}^{\infty} S_2(m,n)\lambda^{-m}\frac{t^m}{m!},\end{aligned} \tag{1.4}$$

where $S_2(m,n)$ is the stirling number of the second kind. When $x=0$, $Ch_{n,\lambda} = Ch_{n,\lambda}(0)$ are called the λ-Changhee numbers. From (1.4), we note that

$$E_m(x) = \sum_{n=0}^{m} Ch_{n,\lambda}(x)S_2(m,n)\lambda^{-m}, \quad (m \geq 0). \tag{1.5}$$

For $n \geq 0$, the stirling number of the first kind is defined as

$$(x)_n = x(x-1)\cdots(x-n+1) = \prod_{l=1}^{n-1}(x-l) = \sum_{l=0}^{n} S_1(n,l)x^l, \tag{1.6}$$

and the stirling number of the second kind is given by

$$x^n = \sum_{l=0}^{n} S_2(n,l)(x)_l, \quad (\text{see } [7,8,9]). \tag{1.7}$$

Let $C(\mathbb{Z}_p)$ be the space of all continuous functions on $\mathbb{Z}_p$. For $f \in C(\mathbb{Z}_p)$, the fermionic p-adic integral on $\mathbb{Z}_p$ is defined by Kim to be

$$\begin{aligned}\int_{\mathbb{Z}_p} f(x)d\mu_{-1}(x) &= \lim_{N\to\infty}\sum_{x=0}^{p^N-1} f(x)\mu_{-1}(x+p^N\mathbb{Z}_p) \\ &= \lim_{N\to\infty}\sum_{x=0}^{p^N-1} f(x)(-1)^x, \quad (\text{see } [8]).\end{aligned} \tag{1.8}$$

From (1.8), we note that

$$I_{-1}(f_1) + I_{-1}(f) = 2f(0), \quad \text{where } f_1(x) = f(x+1). \tag{1.9}$$

Thus, by (1.9), we get

$$I_{-1}(f_n) + (-1)^{n-1}I_{-1}(f) = 2\sum_{l=0}^{n-1}(-1)^{n-1-l}f(l), \quad (\text{see } [8]). \tag{1.10}$$

where $f_n(x) = f(x+n),\ n \in \mathbb{N}$. It is not difficult to show that

$$\begin{aligned}\frac{1-\sqrt{1-4t}}{2t} =& \frac{(-1)}{2t}\left(\sum_{n=1}^{\infty}\binom{2n}{n}\frac{(-1)^{n-1}}{4^n(2n-1)}(-1)^n 4^n t^n\right)\\ =& \frac{1}{2}\sum_{n=1}^{\infty}\binom{2n}{n}\frac{1}{2n-1}t^{n-1}\\ =& \frac{1}{2}\sum_{n=0}^{\infty}\binom{2n+2}{n+1}\frac{1}{2n+1}t^n \qquad (1.11)\\ =& \frac{1}{2}\sum_{n=0}^{\infty}\frac{(2n+2)(2n+1)(2n)\cdots(n+2)}{(n+1)n!(2n+1)}t^n\\ =& \sum_{n=0}^{\infty}\binom{2n}{n}\frac{1}{n+1}t^n, \quad (\text{see } [1,2,3]).\end{aligned}$$

As is well known, the Catalan number C_n is defined by $C_n = \binom{2n}{n}\frac{1}{n+1},\ (n \geq 0)$. From (1.11), we note that the generating function of Catalan numbers is given by

$$\frac{2}{1+\sqrt{1-4t}} = \sum_{n=0}^{\infty} n! C_n \frac{t^n}{n!}, \quad (\text{see } [9-13]). \qquad (1.12)$$

In this paper, we study Catalan numbers associated with p-adic integral on $\mathbb{Z}_p$ and we give Witt's type formula related Catalan numbers.

2. Catalan numbers associated with p-adic intergral on $\mathbb{Z}_p$

For $t \in \mathbb{C}_p$ with $|t|_p < p^{-\frac{1}{p-1}}$, we observe that

$$\int_{\mathbb{Z}_p}(1+t)^{\frac{x}{2}}d\mu_{-1}(x) = \frac{2}{1+\sqrt{1+t}} = \sum_{n=0}^{\infty} Ch_{n,\frac{1}{2}}\frac{t^n}{n!}. \qquad (2.1)$$

From (1.9), we note that

$$\int_{\mathbb{Z}_p} e^{xt}d\mu_{-1}(x) = \frac{2}{e^t+1}\sum_{n=0}^{\infty} E_n \frac{t^n}{n!} \qquad (2.2)$$

On the other hand,

$$\begin{aligned}\int_{\mathbb{Z}_p}(1+t)^{\frac{x}{2}}d\mu_{-1}(x) &= \sum_{m=0}^{\infty}\int_{\mathbb{Z}_p}x^m d\mu_{-1}(x)\frac{1}{2^m}\frac{1}{m!}\Big(\log(1+t)\Big)^m \\ &= \sum_{m=0}^{\infty}E_m 2^{-m}\sum_{n=m}^{\infty}S_1(n,m)\frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty}\left(\sum_{m=0}^{n}2^{-m}E_m S_1(n,m)\right)\frac{t^n}{n!}.\end{aligned} \tag{2.3}$$

Therefore, by (2.1) and (2.3), we obtain the following theorem.

Theorem 1. For $n \geq 0$, we have

$$\sum_{m=0}^{n}2^{-m}E_m S_1(n,m) = Ch_{n,\frac{1}{2}}$$

By replacing t by $-4t$ in (2.1), we get

$$\begin{aligned}\int_{\mathbb{Z}_p}(1-4t)^{\frac{x}{2}}d\mu_{-1}(x) =& \frac{2}{1+\sqrt{1-4t}} = \sum_{n=0}^{\infty}\frac{1}{n+1}\binom{2n}{n}t^n \\ =& \sum_{n=0}^{\infty}C_n t^n.\end{aligned} \tag{2.4}$$

On the other hand,

$$\int_{\mathbb{Z}_p}(1-4t)^{\frac{x}{2}}d\mu_{-1}(x) = \sum_{n=0}^{\infty}Ch_{n,\frac{1}{2}}\frac{(-4)^n t^n}{n!}. \tag{2.5}$$

Therefore, by (2.4) and (2.5), we obtain the following theorem.

Theorem 2. For $n \geq 0$, we have

$$C_n = (-1)^n\frac{4^n Ch_{n,\frac{1}{2}}}{n!}.$$

From (2.4), we have

$$\begin{aligned}\sum_{n=0}^{\infty}(-1)^n 4^n\int_{\mathbb{Z}_p}\binom{\frac{x}{2}}{n}d\mu_{-1}(x)t^n &= \sum_{n=0}^{\infty}C_n t^n \\ &= \sum_{n=0}^{\infty}\frac{1}{n+1}\binom{2n}{n}t^n.\end{aligned} \tag{2.6}$$

Therefore, by comparing the coefficients on the both sides of (2.6), we obtain the following theorem.

Theorem 3. For $n \geq 0$, we have

$$\int_{\mathbb{Z}_p} \binom{\frac{x}{2}}{n} d\mu_{-1}(x) = \frac{(-1)^n}{4^n} C_n = \frac{(-1)^n}{4^n(n+1)} \binom{2n}{n}.$$

From (2.3), we have the following equation:

$$\int_{\mathbb{Z}_p} \binom{\frac{x}{2}}{n} d\mu_{-1}(x) = \frac{1}{n!} \sum_{m=0}^{n} 2^{-m} E_m S_1(n,m) \tag{2.7}$$

Therefore, by Theorem 3 and (2.7), we obtain the following theorem.

Theorem 4. For $n \geq 0$, we have

$$C_n = \frac{(-1)^n}{n!} \sum_{m=0}^{n} 2^{2n-m} E_m S_1(n,m).$$

Now, we observe that

$$\begin{aligned}
\sqrt{1+t} = (1+t)^{\frac{1}{2}} &= \sum_{n=0}^{\infty} \binom{\frac{1}{2}}{n} t^n = \sum_{n=0}^{\infty} \frac{\left(\frac{1}{2}\right)_n}{n!} t^n \\
&= \sum_{n=0}^{\infty} \frac{1\left(\frac{1}{2}-1\right)\left(\frac{1}{2}-2\right)\cdots\left(\frac{1}{2}-n+1\right)}{n!} t^n \\
&= \sum_{n=0}^{\infty} \frac{(-1)^{n-1} 1\cdot 3\cdot 5\cdots(2n-3)}{n! 2^n} t^n \\
&= \sum_{n=0}^{\infty} \frac{(-1)^{n-1} 1\cdot 2\cdot 3\cdot 4\cdots(2n-3)(2n-2)(2n-1)(2n)}{n! 2^n 2\cdot 4\cdot 6\cdots(2n-2)(2n-1)(2n)} t^n \\
&= \sum_{n=0}^{\infty} (-1)^{n-1} \frac{(2n)!}{n! 4^n (2n-1) n!} t^n = \sum_{n=0}^{\infty} \binom{2n}{n} \frac{(-1)^{n-1}}{4^n(2n-1)} t^n.
\end{aligned} \tag{2.8}$$

By (2.1) and (2.8), we get

$$\begin{aligned}
2 &= \left(\sum_{n=0}^{\infty} Ch_{n,\frac{1}{2}}\frac{t^n}{n!}\right)\left(1+\sqrt{1+t}\right)\\
&= \sum_{n=0}^{\infty} Ch_{n,\frac{1}{2}}\frac{t^n}{n!} + \left(\sum_{k=0}^{\infty} Ch_{k,\frac{1}{2}}\frac{t^k}{k!}\right)\left(\sum_{m=0}^{\infty}\binom{2m}{m}\frac{(-1)^{m-1}}{4^m(2m-1)}t^m\right)\\
&= \sum_{n=0}^{\infty} Ch_{n,\frac{1}{2}}\frac{t^n}{n!} + \sum_{n=0}^{\infty}\left(\sum_{m=0}^{n} C_m\frac{(m+1)(-1)^{m-1}}{4^m(2m-1)}\frac{m!n!}{(n-m)!m!}Ch_{n-m,\frac{1}{2}}\right)\frac{t^n}{n!}\\
&= \sum_{n=0}^{\infty} Ch_{n,\frac{1}{2}}\frac{t^n}{n!} + \sum_{n=0}^{\infty}\left(\sum_{m=0}^{n} C_m\frac{(m+1)!(-1)^{m+1}}{4^m(2m-1)}\binom{n}{m}Ch_{n-m,\frac{1}{2}}\right)\frac{t^n}{n!}
\end{aligned} \tag{2.9}$$

By comparing the coefficients on the both sides of (2.9), we get

$$Ch_{n,\frac{1}{2}} + \sum_{m=0}^{n}\binom{n}{m}C_m Ch_{n-m,\frac{1}{2}}(m+1)!(-1)^{m+1}\frac{1}{4^m(2m-1)} = \begin{cases} 2 & \text{if } n=0\\ 0 & \text{if } n>0. \end{cases} \tag{2.10}$$

Therefore, by (2.10), we obtain the following theorem.

Theorem 5. For $n \in \mathbb{N}$, we have

$$Ch_{n,\frac{1}{2}} = \sum_{m=0}^{n}\binom{n}{m}C_m Ch_{n-m,\frac{1}{2}}(-1)^m\frac{(m+1)!}{4^m(2m-1)},$$

and

$$Ch_{0,\frac{1}{2}} = 1.$$

By replacing t by $\frac{1}{4}\left(1-e^{2t}\right)$ in (1.12), we get

$$\begin{aligned}
\frac{2}{e^t+1} &= \sum_{n=0}^{\infty} C_n\frac{1}{4^n}\left(1-e^{2t}\right)^n\\
&= \sum_{n=0}^{\infty} C_n\frac{(-1)^n}{4^n}\left(e^{2t}-1\right)^n\\
&= \sum_{n=0}^{\infty} C_n\frac{(-1)^n}{4^n}n!\sum_{m=n}^{\infty} S_2(m,n)\frac{2^m t^m}{m!}\\
&= \sum_{m=0}^{\infty}\left(\sum_{n=0}^{m} C_n(-1)^n 2^{m-2n}n!S_2(m,n)\right)\frac{t^m}{m!}.
\end{aligned} \tag{2.11}$$

Therefore, by (1.1) and (2.11), we obtain the following theorem.

Theorem 6. For $m \geq 0$, we have

$$E_m = \sum_{n=0}^{m} C_n (-1)^n 2^{m-2n} n! S_2(m,n).$$

Now, we observe

$$\begin{aligned} (1+t)^{\frac{x}{2}} &= \sum_{m=0}^{\infty} \left(\frac{x}{2}\right)^m \frac{\big(\log(1+t)\big)^m}{m!} = \sum_{m=0}^{\infty} \left(\frac{x}{2}\right)^m \sum_{n=m}^{\infty} S_1(n,m) \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} \left(\sum_{m=0}^{n} \left(\frac{x}{2}\right)^m S_1(n,m) \right) \frac{t^n}{n!}. \end{aligned} \tag{2.12}$$

We consider the $\frac{1}{2}$-Changhee polynomials which are given by the generating function to be

$$\begin{aligned} \int_{\mathbb{Z}_p} (1+t)^{\frac{x+y}{2}} d\mu_{-1}(y) &= \frac{2}{1+\sqrt{1+t}} \sqrt{(1+t)^x} \\ &= \sum_{n=0}^{\infty} Ch_{n,\frac{1}{2}}(x) \frac{t^n}{n!}. \end{aligned} \tag{2.13}$$

When $x = 0$, we note that $Ch_{n,\frac{1}{2}} = Ch_{n,\frac{1}{2}}(0)$. By (2.11), (2.12) and (2.13), we get

$$\begin{aligned} \sum_{n=0}^{\infty} Ch_{n,\frac{1}{2}}(x) \frac{t^n}{n!} &= \left(\frac{2}{1+\sqrt{1+t}} \right) \left((1+t)^{\frac{x}{2}} \right) \\ &= \left(\sum_{k=0}^{\infty} Ch_{k,\frac{1}{2}} \frac{t^k}{k!} \right) \left(\sum_{m=0}^{\infty} \left(\sum_{j=0}^{m} \left(\frac{x}{2}\right)^j S_1(m,j) \right) \frac{t^m}{m!} \right) \\ &= \sum_{n=0}^{\infty} \left(\sum_{m=0}^{n} \sum_{j=0}^{m} \left(\frac{x}{2}\right)^j S_1(m,j) Ch_{n-m,\frac{1}{2}} \binom{n}{m} \right) \frac{t^n}{n!}. \end{aligned} \tag{2.14}$$

Therefore, by (2.14), we obtain the following theorem.

Theorem 7. For $n \geq 0$, we have

$$Ch_{n,\frac{1}{2}}(x) = \sum_{m=0}^{n} \sum_{j=0}^{m} \left(\frac{x}{2}\right)^j S_1(m,j) Ch_{n-m,\frac{1}{2}} \binom{n}{m}.$$

By replacing t by $-4t$ in (2.13), we define Catalan polynomials which are given by the generating function to be

$$\begin{aligned}\int_{\mathbb{Z}_p}(1-4t)^{\frac{x+y}{2}}d\mu_{-1}(y) &= \frac{2}{1+\sqrt{1-4t}}\sqrt{(1-4t)^x} \\ &= \sum_{n=0}^{\infty}C_n(x)t^n.\end{aligned} \tag{2.15}$$

From (1.12) and (2.12), we note that

$$\begin{aligned}\sum_{n=0}^{\infty}C_n(x)t^n &= \left(\frac{2}{1+\sqrt{1-4t}}\right)\left((1-4t)^{\frac{x}{2}}\right) \\ &= \left(\sum_{k=0}^{\infty}C_kt^k\right)\left(\sum_{m=0}^{\infty}\left(\sum_{j=0}^{m}\left(\frac{x}{2}\right)^jS_1(m,j)\right)\frac{(-4)^mt^m}{m!}\right) \\ &= \sum_{n=0}^{\infty}\left(\sum_{m=0}^{n}\sum_{j=0}^{m}\left(\frac{x}{2}\right)^jS_1(m,j)\frac{(-4)^m}{m!}C_{n-m}\right)t^n.\end{aligned} \tag{2.16}$$

Therefore, by comparing the coefficients on the both sides of (2.16), we obtain the following theorem.

Theorem 8. For $n \geq 0$, we have

$$C_n(x) = \sum_{m=0}^{n}\sum_{j=0}^{m}\left(\frac{x}{2}\right)^jS_1(m,j)\frac{(-4)^m}{m!}C_{n-m}.$$

Remark. From (2.8) and (2.12), we note that

$$\sum_{n=0}^{\infty}\binom{2n}{n}\frac{(-1)^{n-1}}{4^n(2n-1)}t^n = \sum_{n=0}^{\infty}\left(\frac{1}{n!}\sum_{m=0}^{n}\left(\frac{1}{2}\right)^mS_1(n,m)\right)t^n \tag{2.17}$$

By (2.17), we get

$$\frac{1}{n+1}\binom{2n}{n}\frac{(-1)^{n-1}n+1}{4^n(2n-1)} = \frac{1}{n!}\sum_{m=0}^{n}\left(\frac{1}{2}\right)^mS_1(n,m) \tag{2.18}$$

Thus, from (2.18), we have

$$C_n = \frac{4^n(2n-1)}{(n+1)!}(-1)^{n-1}\sum_{m=0}^{n}\left(\frac{1}{2}\right)^mS_1(n,m),\ (n \geq 0).$$

Corollary 9. For $n \geq 0$, we have

$$C_n = (-1)^{n+1}\frac{4^n(2n-1)}{(n+1)!}\sum_{m=0}^{n}\left(\frac{1}{2}\right)^mS_1(n,m).$$

References

1. H. W. Gould, *Sums and convolved sums of Catalan numbers and their generating functions,* Indian J. Math. 46 (2004), no. 2-3, 137-160.
2. R. Hampel, *On the problem of Catalan,* (Polish) Prace Mat. 4 (1960), 11-19.
3. S. Hyyro, *On the Catalan problem,* (Finnish) Arkhimedes 1963 (1963), no. 1, 53-54.
4. K. Inkeri, *On Catalan's problem,* Acta Arith. 9 (1964), 285-290.
5. D. S. Kim, T. Kim, *Some identities of Korobov-type polynomials associated with p-adic integrals on $\mathbb{Z}_p$,* Advances in Difference Equations (2015) 2015:282 DOI 10.1186/s13662-015-09602-8.
6. T. Kim, D. S. Kim, *A note on nonlinear Changhee differential equations,* Russ. J. Math. Phys. 23 (2016), 88-92.
7. T. Kim D. S. Kim, J.-J. Seo, H.-I. Kwon, *Differential equations associated with λ-Changhee polynomials,* J. Nonlinear Sci. Appl. 9 (2016), 3098-3111.
8. T. Kim, *A study on the q-Euler numbers and the fermionic q-integral of the product of several type q-Bernstein polynomials on $\mathbb{Z}_p$,* Adv. Stud. Contemp. Math. (Kyungshang) 23 (2013), no. 1, 5-11.
9. J. Morgado, *Some remarks on an identity of Catalan concerning the Fibonacci numbers,* Special issue in honor of António Monteiro. Portugal. Math. 39 (1980), no. 1-4, 341-348 (1985).
10. A. Natucci, *Ricerche sistematiche intorno al "teorema di Catalan",* (Italian) Giorn. Mat. Battaglini. (5) 2 (82) (1954), 297-300.
11. R. Rangarajan, P. Shashikala, *A pair of classical orthogonal polynomials connected to Catalan numbers,* Adv. Stud Contemp. Math. (Kyungshang) 23 (2013), no. 2, 323-335.
12. A. Rotkiewicz, *Sur le probléme de Catalan,* (French) Elem. Math. 15 (1960), 121-134.
13. A. D. Sands, *On generalised Catalan numbers,* Discrete Math. 21 (1978), no. 2, 219-221.

Department of Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea

E-mail address: tkkim@kw.ac.kr